

HR: 12:05h

AN: **V42B-08** [Abstracts]

TI: **Crustal recycling model: testing by linking petrology with isotope geochemistry**

AU: \* **Sobolev, A V**

EM: *asobolev@mpch-mainz.mpg.de*

AF: *Max-Planck Institute for Chemistry, P.O. Box 3060, Mainz, 55020, Germany*

AU: \* **Sobolev, A V**

EM: *asobolev@mpch-mainz.mpg.de*

AF: *Vernadsky Institute of Geochemistry, Russian Academy of Sciences, Kosygin street 19, Moscow, 119991, Russian Federation*

AU: **Hofmann, A W**

EM: *hofmann@mpch-mainz.mpg.de*

AF: *Max-Planck Institute for Chemistry, P.O. Box 3060, Mainz, 55020, Germany*

AU: **Brueggemann, G**

EM: *brueggemag@uni-mainz.de*

AF: *Institut für Geowissenschaften, Johannes Gutenberg-Universität Mainz,, Becherweg 21,, Mainz, 55129, Germany*

AU: **Batanova, V G**

EM: *batanova@mpch-mainz.mpg.de*

AF: *Vernadsky Institute of Geochemistry, Russian Academy of Sciences, Kosygin street 19, Moscow, 119991, Russian Federation*

AU: **Sigurdsson, I A**

EM: *ingvar@nattsud.is*

AF: *South Iceland Nature Centre, Strandvegur 50, Vestmannaeyjar, IS 900, Iceland*

AU: **Kuzmin, D V**

EM: *kuzmin@mpch-mainz.mpg.de*

AF: *V.S. Sobolev Institute of Geology and Mineralogy, Siberina Branch of Russian Academy of Sciences, Koptuyga prosp. 3, Novosibirsk, 630090, Russian Federation*

AU: **Kuzmin, D V**

EM: *kuzmin@mpch-mainz.mpg.de*

AF: *Max-Planck Institute for Chemistry, P.O. Box 3060, Mainz, 55020, Germany*

AB: Recycling of subducted oceanic crust is widely thought to explain much of the chemical and isotopic heterogeneity of Earth's present-day mantle (1). Recycled subducted ocean crust has been traced by elevated 187Os/188Os in some studies (2-4) and by high Ni and low Mn contents in others (5,6). Recently, we have linked these tracers for the first time (7). We combine data for averaged compositions of olivine phenocrysts for olivine rich lavas and bulk rock 187Os/188Os ratios from Iceland and Hawaii, and we apply the method (6,7) to obtain the proportion of pyroxenite derived melt by both Ni excess and Mn deficit in olivine. The observed strong correlation for quaternary lavas of Iceland significantly strengthens the recycling model (1,8). It allows us to estimate the Os isotopic composition of both the recycled crust and the mantle peridotite, thereby constraining the model ages of end-members. We show that Icelandic quaternary lavas require an ancient crustal component with model ages between 1.1 and 1.8 Ga and a peridotite end-member close to present-day oceanic mantle. The similar data for Hawaii suggest younger ages of recycled material: 0.5-1 Ga. References: 1. A.W. Hofmann, W. M. White, EPSL. 57, 421 (1982) 2. J. C. Lassiter, E. H. Hauri, EPSL 164, 483 (1998). 3. A.C. Skovgaard et al, EPSL 194, 259 (2001). 4. A. D. Brandon et al., GCA

71, 4570 (2007). 5. A.V. Sobolev et al., Nature 434, 590 (2005). 6. A.V. Sobolev et al., Science 316, 412 (2007). 7. A.V. Sobolev et al., Science 321, 536 (2008). 8. C. Hemond et al., JGR–Solid Earth 98, 15833 (1993).

DE: 1025 Composition of the mantle

DE: 1037 Magma genesis and partial melting (3619)

DE: 1038 Mantle processes (3621)

DE: 1065 Major and trace element geochemistry

DE: 3640 Igneous petrology

SC: Volcanology, Geochemistry, Petrology [V]

MN: 2008 Fall Meeting